Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 1 week, comes back, or occurs with a fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

236v2.1

PEEL

Double Strength Expectorant DM Cough Syrup

Drug Facts

Active ingredients   Purposes
(in each teaspoonful (5 mL))

Dextromethorphan HBr 20 mg . . . . Cough suppressant
Guaifenesin 300 mg . . . . Expectorant

Uses
- helps loosen phlegm (mucus) and thin bronchial secretions to:
  - drain bronchial tubes
  - make coughs more productive
  - rid the bronchial passageways of bothersome mucus
- temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ▶

Drug Facts (continued)

Directions
- take every 4 hours
- do not take more than 6 doses in 24 hours

adults and children 12 years and over	1 teaspoonful (5 mL)
children 6 to under 12 years	1/2 teaspoonful (2-1/2 mL)
children under 6 years	ask a doctor

Other information
- store between 15 ˚C and 30 ˚C (59 ˚F and 86 ˚F)

Inactive ingredients
citric acid, flavor, menthol, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

Weeks & Leo Co., Inc. Tel: (515) 276-1586