

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

PEEL

Sinutrol PE Tablets

Drug Facts

Active ingredients (in each tablet)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves these cold symptoms:

nasal congestion	sinus congestion and pressure	
runny nose	minor aches and pains	headache
sneezing	temporarily reduces fever	

temporarily relieves these symptoms of hay fever or other upper respiratory allergies:

runny nose	sneezing	
nasal congestion	headache	itchy, watery eyes
sinus congestion and pressure	itching of the nose or throat	
minor aches and pains		

23742.1

Outside label

Drug Facts (continued)

Ask a doctor before use if you have

glaucoma	diabetes	heart disease	thyroid disease
high blood pressure		liver disease	
a breathing problem such as emphysema or chronic bronchitis			
trouble urinating due to an enlarged prostate gland			

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers
taking the blood thinning drug warfarin

When using this product

do not use more than directed

drowsiness may occur
avoid alcoholic drinks
alcohol, sedatives, and tranquilizers may increase drowsiness
excitability may occur, especially in children
be careful when driving a motor vehicle or operating machinery

Inside label

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 12 tablets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

Stop use and ask a doctor if you get nervous, dizzy, or sleepless
redness or swelling is present new symptoms occur
fever gets worse or lasts more than 3 days
pain or nasal congestion gets worse or lasts more than 7 days or is accompanied by fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over:
take 2 tablets every 4 hours
do not take more than 12 tablets in 24 hours
children under 12 years of age: ask a doctor

Drug Facts (continued)

Other information

store at room temperature 15 °C to 25 °C (59 °F to 77 °F)
to report serious side effects due to the use of this product call
(515) 276-1586

Inactive ingredients

corn starch, croscarmellose sodium, D&C yellow no. 10 lake, FD&C yellow no. 6 lake, hypromellose, magnesium stearate, microcrystalline cellulose, silica, sodium starch glycolate, stearic acid

Base label - adheres to bottle