

Tamper-evident: do not use if the imprinted seal under the cap is torn, broken, or missing.

Diphenhydramine HCl Caplets 25 mg

Drug Facts

Active ingredient (in each caplet)	Purpose
Diphenhydramine HCl 25 mg	Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing

137/2,4

PEEL

Outside label

Drug Facts (continued)

Directions

- take every 4 to 6 hours
 - do not take more than 6 doses in 24 hours
- | | |
|---------------------------------------|----------------|
| adults and children 12 years and over | 1 or 2 caplets |
| children 6 years to under 12 years | 1 caplet |
| children under 6 years | ask a doctor |

Other information

- each tablet contains: calcium 25 mg
- protect from light
- store at 25 °C (77 °F) excursions permitted between 15 °C and 30 °C (59 °F and 86 °F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C red no. 27 aluminum lake, dibasic calcium phosphate dihydrate, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polysorbate 80, titanium dioxide

To report serious side effects call 515-276-1586.

Base label

Drug Facts (continued)

Warnings

Do not use with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Inside label