

PEEL

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

Loratadine Tablets 10 mg

Drug Facts	
Active ingredient (in each tablet)	Purpose
Loratadine USP, 10 mg	Antihistamine
Uses	
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:	
runny nose	
itchy, watery eyes	
sneezing	
itching of the nose or throat	
Warnings	
Do not use if you have ever had an allergic reaction to this product or any of its ingredients	

232v2.0

Outside label

Drug Facts (continued)

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at **1-800-FDA-1088**.
an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Inside label

Drug Facts (continued)

Directions

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with kidney or liver disease	ask a doctor

Other information
store between 20 and 25 °C (68 and 77 °F)
protect from excessive moisture

Inactive ingredients
corn starch, lactose monohydrate, magnesium stearate, pregelatinized starch

Base label - adheres to bottle