

Extra Strength Super Troche DM

Drug Facts

Active ingredients (in each lozenge)

Purposes

Benzocaine 15 mg Oral pain reliever
Dextromethorphan HBr 10 mg Cough suppressant

Uses temporarily relieves these cold symptoms:
■ cough ■ minor sore throat

Warnings

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Sore throat warning: if sore throat is severe, lasts for more than 2 days, occurs with or followed by fever, headache, rash, swelling, nausea or vomiting, consult a doctor right away.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with a fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 6 years and over: allow 1 lozenge to dissolve slowly in the mouth. Repeat every 4 hours as needed or as directed by a doctor.
- children under 6 years of age: ask a doctor.

Other information ■ store in a cool, dry place

Inactive ingredients dextrose, FD&C blue no. 1, FD&C red no. 40, flavor, glucose, glycerin, propylene glycol, sucrose, vanillin