

**PEEL**

**Tamper-evident:** do not use if the imprinted seal under the cap is broken or missing.

**Pain & Sleep Caplets**

**Drug Facts**

Active ingredients (in each caplet)	Purposes
Acetaminophen 500 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg .....	Nighttime sleep-aid

**Uses** temporarily relieves minor aches and pains and accompanying sleeplessness due to:  
 headache      muscular aches      minor pain from arthritis  
 backache      premenstrual and menstrual cramps

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 2 caplets in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product

1659/2.5

Outside label

**Drug Facts (continued)**

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. with any other product containing diphenhydramine, even one used on skin for children under 12 years of age

**Ask a doctor before use if you have** liver disease      glaucoma  
 a breathing problem such as emphysema or chronic bronchitis  
 trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin      taking sedatives or tranquilizers

**When using this product** avoid alcoholic drinks      drowsiness will occur      do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if**  sleeplessness lasts continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.      pain gets worse or lasts more than 10 days  
 fever gets worse or lasts more than 3 days      redness or swelling is present      any new symptoms appear

Inside label

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

**Directions** do not take more than recommended adults and children 12 years and over: take 2 caplets at bedtime if needed, or as directed by a doctor      children under 12 years: ask a doctor

**Other information** store between 20 °C and 25 °C (68 °F and 77 °F) to report serious side effects due to the use of this product call (515) 276-1586

**Inactive ingredients**  
 colloidal silicon dioxide, croscarmellose sodium, FD&C blue no. 1 aluminum lake, microcrystalline cellulose, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Base label - adheres to bottle