

Tamper-evident: do not use if either the imprinted seal around the cap or the imprinted seal under the cap is broken or missing.

G-Fenesin DM Caplets

Drug Facts

Active ingredients (in each caplet) Purposes

Dextromethorphan HBr 20 mgCough suppressant
Guaifenesin 400 mgExpectorant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to:
 - drain bronchial tubes
 - make coughs more productive
 - rid the bronchial passageways of bothersome mucus
- temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

238v2.0

Lift Here →

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or occurs with a fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. →

Hinge

Drug Facts (continued)

Directions

- take every 4 hours
- do not take more than 6 doses in 24 hours

adults and children 12 years and over	1 caplet
children 6 years to under 12 years	1/2 caplet
children under 6 years	ask a doctor

Other information

store at 15 °C to 30 °C (59 °F to 86 °F)

Inactive ingredients

colloidal silicon dioxide, magnesium stearate, microcrystalline cellulose, maltodextrin, povidone, stearic acid

Hinge