

Drug Facts (continued)

Other information

- read the directions, warnings, and package insert before use ■ keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity ■ protect product from moisture

Inactive ingredients carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

0 257-43A3.0

42
 Tablets
 Three 14-Day
 Courses of Treatment

Acid Reducer

Delayed Release Tablets 20 mg

Omeprazole

Treats Frequent Heartburn!
 occurring 2 or more days a week



Compare to Prilosec® OTC
 active ingredient only

Item # 12042 (42 Tablets)



Distributed by:
 Weeks & Leo Co., Inc.
 4000 NW 100th St
 Des Moines, IA 50322

Batch #

Exp. Date

Tamper-evident: do not use if any blister unit is torn, broken, or shows any signs of tampering.

Your
 Hometown
 Pharmacy!

Drug Facts

Active ingredient (in each tablet)

Omeprazole delayed-release tablet, 20 mg Acid reducer

Purpose

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

■ frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking ■ warfarin or clopidogrel (blood-thinning medicine) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus (immune system medicine) ■ prescription antiretrovirals (medicines for HIV infection)

Stop use and ask a doctor if ■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088 ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- 14-Day Course of Treatment**
 - swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day
 - do not chew or crush the tablets ■ do not crush tablets in food ■ do not use for more than 14 days unless directed by your doctor
- Repeated 14-Day Courses (if needed)**
 - you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Glue Flap

Glue Flap

Glue Flap