

PEEL

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

Stool Softener Softgels 100 mg

Drug Facts

Active ingredient (in each softgel)	Purpose
Docusate sodium 100 mg.....	Stool softener laxative

Uses
for the relief of occasional constipation (irregularity)
generally produces bowel movement in 12 to 72 hours

Warnings
Ask a doctor before use if you have
stomach pain nausea vomiting
noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Ask a doctor or pharmacist before use if you are presently taking mineral oil ▶

01862.1

Outside label

Drug Facts (continued)

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition you need to use a laxative longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
take softgels with a full glass (8 oz) of water

adults and children 12 years and over	take 1 to 3 softgels daily. This dose may be taken as a single daily dose or in divided doses.
children 2 to under 12 years	take 1 softgel daily
children under 2 years	ask a doctor ▶

Inside label

Drug Facts (continued)

Other information
each softgel contains: sodium 5 mg
store at room temperature between 15 °C and 30 °C (59 °F and 86 °F)
to report serious side effects due to the use of this product call (515) 276-1586

Inactive ingredients
edible ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol 400, propylene glycol, sorbitol

Base label - adheres to bottle