

Tamper-evident: this product protected with sealed blister units. Do not use if any are torn or broken.

Ultra A-D Caplets

Drug Facts

Active ingredient (in each caplet)	Purpose
Loperamide HCl USP, 2 mg	Anti-diarrheal

Use
controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

160V4.2

Lift Here →

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- diarrhea lasts for more than 2 days
- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on next page. If possible, use weight to dose; otherwise, use age.

Hinge

Drug Facts (continued)

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 years to 11 years (60 lbs to 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 years to 8 years (48 lbs to 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information store between 15 °C and 30 °C (59 °F and 86 °F)

Inactive ingredients anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10, FD&C blue no. 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Hinge