

**Tamper-evident:** this product protected with sealed blister units. Do not use if any are torn or broken.

### Ultra A-D Caplets

#### Drug Facts

Active ingredient (in each caplet)	Purpose
Loperamide HCl USP, 2 mg .....	Anti-diarrheal

#### Use

controls symptoms of diarrhea, including Travelers' Diarrhea

#### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use** if you have bloody or black stool

#### Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

160V4.2

Lift Here →

→

→

#### Drug Facts (continued)

**Ask a doctor or pharmacist before use if you are** taking antibiotics

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

#### Stop use and ask a doctor if

- diarrhea lasts for more than 2 days
- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on next page. If possible, use weight to dose; otherwise, use age.

Hinge

#### Drug Facts (continued)

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 years to 11 years (60 lbs to 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 years to 8 years (48 lbs to 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

**Other information** store between 15 °C and 30 °C (59 °F and 86 °F)

**Inactive ingredients** anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10, FD&C blue no. 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Hinge