

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

Acetaminophen Caplets 500 mg

Drug Facts	
<i>Active ingredient (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg.....	Pain reliever/fever reducer

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - minor pain from arthritis
 - headache
 - backache
 - muscular aches
 - a cold
 - flu
 - premenstrual and menstrual cramps
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Lift Here

138x2.4

Outside label

Drug Facts (continued)

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure if a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Inside label

Drug Facts (continued)

Directions

- adults and children 12 years and over: take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at room temperature between 15 °C and 30 °C (59 °F and 86 °F)
- call **515-276-1586** to report serious side effects due to the use of this product

Inactive ingredients

hypromellose, polyethylene glycol, povidone, pregelatinized corn starch, stearic acid, titanium dioxide. Also contains carboxymethylcellulose sodium or sodium starch glycolate.

Base label - adheres to bottle