

**Tamper-evident:** do not use if the imprinted seal under the cap is broken or missing.

### Acetaminophen Caplets 500 mg

#### Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever/fever reducer

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - minor pain from arthritis
  - muscular aches
  - premenstrual and menstrual cramps
  - headache
  - a cold
  - toothache
  - backache
  - flu

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Lift Here

138x2.4

Outside label

#### Drug Facts (continued)

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure if a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Drug Facts (continued)

#### Directions

- adults and children 12 years and over: take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

#### Other information

- store at room temperature between 15 °C and 30 °C (59 °F and 86 °F)
- call **515-276-1586** to report serious side effects due to the use of this product

#### Inactive ingredients

hypromellose, polyethylene glycol, povidone, pregelatinized corn starch, stearic acid, titanium dioxide. Also contains carboxymethylcellulose sodium or sodium starch glycolate.

Inside label

Base label - adheres to bottle