

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

Ibuprofen Caplets 200 mg

Drug Facts

Active ingredient (in each caplet)	Purpose
Ibuprofen, USP 200 mg (NSAID)*.....	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily relieves minor aches and pains due to:

headache	the common cold	muscular aches
backache	menstrual cramps	toothache
minor pain of arthritis		
temporarily reduces fever		

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives	facial swelling	asthama (wheezing)	shock
skin reddening	rash	blisters	

15112,5

Outside label

Drug Facts (continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have problems or serious side effects from taking pain relievers or fever reducers

Inside label 1

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug

When using this product take with food or milk if stomach upset occurs long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at **1-800-FDA-1088** you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is

Base label - 1

Drug Facts (continued)

especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed the smallest effective dose should be used

do not take longer than 10 days, unless directed by a doctor (see Warnings)

adults and children 12 years and older	take one caplet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 caplet, 2 caplets may be used do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Inside label 2

Drug Facts (continued)

Other information

store between 20-25 °C (68-77 °F)

read all directions and warnings before use

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, magnesium stearate, microcrystalline, cellulose, polydextrose, polyethylene glycol, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Base label - adheres to bottle