

Tamper evident: do not use if the imprinted seal under the cap is broken or missing.

Peel

Naproxen Sodium Tablets 220 mg

Drug Facts

Active ingredient (in each tablet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily relieves minor aches and pains due to:

minor pain of arthritis	backache	headache
the common cold	muscular aches	
menstrual cramps	toothache	

temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives	asthma (wheezing)	skin reddening
-------	-------------------	----------------

221V2.4

Outside label

Drug Facts (continued)

facial swelling shock rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic

Inside label 1

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking any other drug containing an NSAID (prescription or nonprescription) taking a blood thinning (anticoagulant) or steroid drug under a doctor's care for any serious condition taking any other drug

When using this product take with food or milk if stomach upset occurs long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if side effects occur. You may report side effects to FDA at **1-800-FDA-1088**.

you experience any of the following signs of stomach bleeding:

feel faint	vomit blood	have bloody or black stools
have stomach pain that does not get better		
pain gets worse or lasts more than 10 days	fever gets worse or lasts more than 3 days	you have difficulty swallowing
like the pill is stuck in your throat	you develop heartburn	stomach pain or upset gets worse or lasts
redness or swelling is present in the painful area	any new symptoms appear	

Base label - 1

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed
the smallest effective dose should be used
do not take longer than 10 days, unless directed by a doctor (see Warnings)
drink a full glass of water with each dose

adults and children 12 years and older	take one tablet every 8 to 12 hours while symptoms last for the first dose you may take 2 tablets within the first hour do not exceed 2 tablets in any 8- to 12-hour period do not exceed 3 tablets in a 24-hour period
--	---

Inside label 2

Drug Facts (continued)

children under 12 years	ask a doctor
-------------------------	--------------

Other information

each tablet contains: sodium 20 mg
store at 20-25 °C (68-77 °F). Avoid high humidity and excessive heat above 40 °C (104 °F)

Inactive ingredients

FD&C Blue #2, croscarmellose sodium, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

Base label - adheres to bottle