

**Tamper-evident:** do not use if the imprinted seal under the cap is broken or missing.

**Aspirin Enteric Coated Tablets 81 mg  
Adult Low Strength**

**Drug Facts**

| <b>Active ingredient (in each tablet)</b> | <b>Purpose</b>              |
|-------------------------------------------|-----------------------------|
| Aspirin 81 mg (NSAID*) .....              | Pain reliever/fever reducer |
| *nonsteroidal anti-inflammatory drug      |                             |

**Uses**

- temporarily relieves minor aches and pains
- temporarily reduces fever
- ask your doctor about other uses for this low strength aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use

1586.2

Lift Here

Outside label (Panel 1)

**Drug Facts (continued)**

- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for diabetes, gout, or arthritis

Inside label (Panel 3)

**Drug Facts (continued)**

this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks everyday while using this product

Inside label (Panel 2)

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - any new symptoms appear
  - redness or swelling is present
  - ringing in the ears or loss of hearing occurs
- have bloody or black stools
- have stomach pain that does not get better

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Inside label (Panel 4)

**Drug Facts (continued)**

**Directions**

- adults and children 12 years and over:
  - drink a full glass of water with each dose every 4 hours while symptoms last
  - take 4 to 8 tablets every 4 hours, unless directed by a doctor
  - do not take more than 48 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor
- ask your doctor about recommended doses for other uses

**Other information**

- store at controlled room temperature; 15 °C to 30 °C (59 °F to 86 °F)
- to report serious side effects due to the use of this product call (515) 276-1586

**Inactive ingredients**

acetylated monoglycerides, anhydrous lactose, colloidal silicon dioxide, croscarmellose sodium, D&C yellow no. 10, hypromellose, hypromellose phthalate, iron oxide ochre, microcrystalline cellulose, mineral oil, polyethylene glycol 400, polysorbate 80, titanium dioxide

Attaches to bottle (Panel 5)