

Tamper-evident: do not use if the imprinted seal under the cap is broken or missing.

Aspirin Enteric Coated Tablets 325 mg

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID*)	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscular aches
 - a cold
 - minor pain from arthritis
 - premenstrual and menstrual cramps
- temporarily reduces fever

05012.2

Lift Here

Outside label (Panel 1)

Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis

Inside label (Panel 3)

Drug Facts (continued)

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

Inside label (Panel 2)

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - any new symptoms appear
 - redness or swelling is present
 - ringing in the ears or loss of hearing occurs
- have bloody or black stools
- have stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Inside label (Panel 4)

Drug Facts (continued)

Directions

- adults and children 12 years and over:
 - drink a full glass of water with each dose
 - take 1 or 2 tablets every 4 hours while symptoms last
 - do not take more than 12 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at controlled room temperature 15 °C to 30 °C (59 °F to 86 °F)
- report serious side effects due to the use of this product call (515) 276-1586

Inactive ingredients

acetylated monoglycerides, corn starch, croscarmellose sodium, D&C yellow no. 10, FD&C yellow no. 6, hypromellose, hypromellose phthalate, microcrystalline cellulose, mineral oil, polyethylene glycol 400, polysorbate 80, titanium dioxide

Attaches to bottle (Panel 5)